



‘সমানো মন্ত্র: সমিতি: সমানী’

**UNIVERSITY OF NORTH BENGAL**  
B.Sc. Honours 6th Semester Examination, 2023

**CC13-MATHEMATICS****RING THEORY AND LINEAR ALGEBRA-II**

Time Allotted: 2 Hours

Full Marks: 60

*The figures in the margin indicate full marks.***GROUP-A****Answer any four questions from the following**

3×4 = 12

1. Show that the polynomial  $3x^5 + 15x^4 - 20x^3 + 10x + 20$  is irreducible over  $\mathbb{Q}$ .
2. Suppose that  $a, b$  are two elements in an integral domain,  $b \neq 0$  and  $a$  is not a unit. Show that  $\langle ab \rangle$  is contained in  $\langle b \rangle$ .
3. Let  $V = C[0, 1]$  and define  $\langle f, g \rangle = \int_0^{1/2} f(t)g(t) dt$ . Is this an inner product on  $V$ ?
4. Prove that the ideal  $\langle x^2 + 1 \rangle$  is prime in  $\mathbb{Z}[x]$  but not maximal in  $\mathbb{Z}[x]$ .
5. Let  $V = P_1(\mathbb{R})$  and for  $p(x) \in V$ , define  $f_1, f_2 \in V^*$ , by  $f_1(p(x)) = \int_0^1 p(t) dt$  and  $f_2(p(x)) = \int_0^2 p(t) dt$ . Prove that  $\{f_1, f_2\}$  is a basis for  $V^*$ .
6. Prove that  $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$  is not diagonalizable.

**GROUP-B****Answer any four questions from the following**

6×4 = 24

7. (a) Prove that every Euclidean Domain is a PID. 4
- (b) Prove that in an integral domain, associates of every irreducible element are also irreducible. 2
8. (a) Let  $T$  be a linear operator on a finite dimensional vector space  $V$  and let  $W$  be a  $T$ -invariant subspace of  $V$ . Then prove that the characteristic polynomial of  $T_W$  divides the characteristic polynomial of  $T$ . 3

- (b) Let  $V$  be an inner product space and  $S_1$  and  $S_2$  be two subsets of  $V$ . Then prove that  $S_1 \subseteq S_2 \Rightarrow S_1^\perp \subseteq S_2^\perp$ . 3
9. (a) Let  $F$  be a field and  $p(x) \in F[x]$ . Then prove that  $\langle p(x) \rangle$  is a maximal ideal in  $F[x]$  iff  $p(x)$  is irreducible over  $F$ . 4
- (b) Prove that 2 and 5 are not irreducible elements in  $\mathbb{Z}[i]$ . 2
- 10.(a) Prove that in an integral domain, two elements  $a$  and  $b$  are associates iff  $\langle a \rangle = \langle b \rangle$ . 3
- (b) Show that  $1 + 2i$  and  $3 + 5i$  are prime to each other in  $\mathbb{Z}[i]$ . 3
- 11.(a) In  $\mathbb{R}^3$ , with standard inner product, let  $P$  be the subspace  $\text{span}\{(1, 1, 0), (0, 1, 1)\}$ . Find  $P^\perp$ . 3
- (b) Let  $V$  be a finite dimensional Euclidean space. Then prove that a linear mapping  $T : V \rightarrow V$  is orthogonal iff  $T$  maps an orthonormal basis to another orthonormal basis. 3
- 12.(a) Prove that the set of all normal operators is a closed subset of  $L(H, H)$  which contains the set of all self-disjoint operators. 4
- (b) Suppose  $A \in L(H, H)$ . Then prove that  $\langle Ax, x \rangle = 0$  for all  $x \in H$  iff  $A = 0$ . 2

**GROUP-C**

**Answer any two questions from the following**

12×2 = 24

- 13.(a) Let  $A : H \rightarrow H$  is a continuous linear operator, where  $H$  is a Hilbert space. Prove that  $A^*$  is a continuous linear operator with  $\|A^*\| = \|A\|$ . 5
- (b) Prove that the dual space of an  $n$  dimensional vector space is  $n$  dimensional. 4
- (c) Find the minimal polynomial of the matrix 3

$$A = \begin{pmatrix} -3 & 2 \\ 0 & -3 \end{pmatrix}$$

- 14.(a) Let  $T$  be an linear operator on  $V = \mathbb{R}^2$  defined by  $T(a, b) = (2a - 2b, -2a + 5b)$  for all  $(a, b) \in \mathbb{R}^2$ . Determine whether  $T$  is normal, self-adjoint or neither. Produce an orthonormal basis of eigenvectors of  $T$  for  $V$ . 6
- (b) Let  $V = P_1(\mathbb{R})$  and  $W = \mathbb{R}^2$  with respective standard ordered bases  $\beta$  and  $\gamma$ . Define  $T : V \rightarrow W$  by  $T(p(x)) = (p(0) - 2p(1), p(0) + p'(0))$ ; where  $p'(x)$  denotes the derivative of  $p(x)$ . Then compute  $[T^t]_{\gamma^*}^{\beta}$  and  $[T]_{\beta}^{\gamma}$ . 6

- 15.(a) Apply Gram-Schmidt process to the subset  $S = \{1, x, x^2\}$  of the inner product space  $V = P_2(\mathbb{R})$  with inner product 4

$$\langle f, g \rangle = \int_0^1 f(t)g(t) dt$$

to obtain an orthonormal basis  $\beta$  for  $\text{span}(S)$ .

- (b) For two subspaces  $W_1$  and  $W_2$  of a finite dimensional vector space  $V$ , prove that 3

$$(W_1 + W_2)^0 = W_1^0 \cap W_2^0.$$

- (c) Let  $T$  be a linear operator on a finite dimensional vector space  $V$  and let  $f(t)$  be the characteristic polynomial of  $T$ . Then prove that  $f(T) = T_0$ , the zero transformation. 5

- 16.(a) Show that the following polynomials are irreducible: 6

(i)  $x^3 - [9]$  over  $\mathbb{Z}_{11}$ .

(ii)  $x^4 + 2x + 2$  over  $\mathbb{Q}$ .

(iii)  $x^6 + x^3 + 1$  over  $\mathbb{Q}$ .

- (b) Let  $R$  be the ring  $\mathbb{Z} \times \mathbb{Z}$ . Show that the linear equation  $(5, 0)x + (20, 0) = (0, 0)$  has infinitely many roots in  $R$ . 3

- (c) In  $\mathbb{Z}_7[x]$ , factorize  $f(x) = x^3 + [1]$  into linear factors. 3

—x—