



'সমানো মন্ত্র: সমিতি: সমানী'

UNIVERSITY OF NORTH BENGAL
B.Sc. Honours 6th Semester Examination, 2023

DSE-P4-MATHEMATICS

Time Allotted: 2 Hours

Full Marks: 60

*The figures in the margin indicate full marks.
Symbols have their usual meanings.*

The question paper contains DSE-4A and DSE-4B. Candidates are required to answer any one from the two courses and they should mention it clearly on the Answer Book.

DSE-4A

DIFFERENTIAL GEOMETRY

GROUP-A

Answer any four questions from the following

3×4 = 12

1. If $\vec{r} = \vec{r}(s)$ is the position vector of a point P with arc-length as parameter on a curve- γ , then show that $\kappa^2 \tau = [\vec{r}', \vec{r}'', \vec{r}''']$.
2. Obtain the equation of the circular helix $r = (a \cos u, a \sin u, bu)$, $-\infty < u < \infty$, where $a > 0$, referred to s as parameter.
3. Determine $f(u)$ so that the curve $r = (a \cos u, a \sin u, f(u))$ shall be a plane.
4. Find the involutes of a helix.
5. Calculate the torsion of the cubic curve $r = (u, u^2, u^3)$.
6. Show that the surface $e^z \cos x = \cos y$ is minimal.

GROUP-B

Answer any four questions from the following

6×4 = 24

7. Show that the radius of curvature ρ and radius of torsion σ of the curve $r = (a \cos u, a \sin u, a \cos 2u)$ at $u = \frac{\pi}{4}$ are $\rho = \frac{5a}{4}$ and $\sigma = \frac{5a}{6}$. 3+3
8. (a) Find the equation of the osculating plane at a point u of the curve $r = (a \cos u, a \sin u, bu)$ 3
- (b) Find the Serret-Frenet approximation of the curve $r = (\cos u, \sin u, u)$ at $u = \frac{\pi}{2}$. 3

9. Prove that the tangent plane to the surface $xyz = a^3$ and the coordinate planes bound a constant volume. 6
10. Define developable surface. Prove that a surface is developable if and only if the specific curvature is zero at all points. 1+5
11. Find the curvature and torsion of the locus of centre of spherical curvature. 3+3
- 12.(a) Prove that the asymptotic lines are orthogonal if and only if the surface is minimal. 3
- (b) Show that the parametric curve on the surface $(u \cos v, u \sin v, v)$ are asymptotic lines. 3

GROUP-C

Answer any *two* questions

12×2 = 24

- 13.(a) Show that the necessary and sufficient condition that a curve be a helix is that 6

$$[r^{IV}, r''', r''] = -\kappa^5 \frac{d}{ds} \left(\frac{\tau}{\kappa} \right) = 0.$$
- (b) Prove that the geodesic curvature of a geodesic on a surface is zero and conversely. 6
14. Prove that the shortest distance between the principal normal at two consecutive points on a curve is 6+6

$$\frac{\rho ds}{\sqrt{\rho^2 + \sigma^2}}$$
 and the line of this distance divides the radius of curvature in the ratio $\rho^2 : \sigma^2$.
- 15.(a) Define first fundamental form. Prove that the first fundamental form is invariant under a transformation of parameters. 1+5
- (b) Show that the curve $r = r(s)$ is asymptotic line if only if $\frac{dr}{ds} \cdot \frac{dN}{ds} = 0$, where N is the surface normal. Write the necessary and sufficient condition for a curve to be a geodesic. 4+2
- 16.(a) Prove that the curves of the family $\frac{v^3}{u^2} = \text{constant}$ are geodesics on a surface 6
 with the metric $ds^2 = v^2 du^2 - 2uv du dv + 2u^2 dv^2, u > 0, v > 0$.
- (b) Define normal curvature. Find the normal curvature of the right angular helicoid $r(u, v) = (u \cos v, u \sin v, cv)$ at a point on it. 1+5

DSE-4B
THEORY OF EQUATIONS

GROUP-A

Answer any four questions

3×4 = 12

1. If α, β, γ are the roots of the equation $x^3 + px^2 + qx + r = 0$, then find the value of $(\alpha + \beta)(\beta + \gamma)(\gamma + \alpha)$. 3
2. Find the remainder when $2x^4 - 6x^3 + 7x^2 - 5x + 1$ is divided by $(2x - 3)$. 3
3. Solve the equation $4x^4 - 4x^3 - 13x^2 + 9x + 9 = 0$ given that the sum of two roots is zero. 3
4. Reduce the reciprocal equation $x^5 - 6x^4 + 7x^3 + 7x^2 - 6x + 1 = 0$ to its standard form. 3
5. Apply Descartes' rule of sign to determine the nature of the roots of the equation $x^{10} - 1 = 0$. 3
6. Obtain the equation whose roots are twice the roots of the equation $x^3 + 3x^2 + 4x + 5 = 0$. 3

GROUP-B

Answer any four questions

6×4 = 24

7. If α be an imaginary root of the equation $x^7 - 1 = 0$, find the equation whose roots are $\alpha + \alpha^6, \alpha^2 + \alpha^5, \alpha^3 + \alpha^4$. 6
8. Solve the following equation by Ferrari's method: 6

$$x^4 + 12x - 5 = 0$$
9. Apply Sturm's theorem to prove that the equation $x^3 - 7x + 7 = 0$ has two roots lying between 1 and 2, and one root lying between -4 and -3. 6
- 10.(a) Show that the equation $2x^7 + 3x^4 + 3x + k = 0$ has at least four complex roots for all values of k . 2
- (b) If α is a root of the equation $x^4 + px^3 - 6x^2 - px + 1 = 0$, then prove that $\frac{1 + \alpha}{1 - \alpha}$ is also a root of it. 4
- 11.(a) Find the multiple roots of the equation 4

$$x^4 - 2x^3 - 11x^2 + 12x + 36 = 0.$$
- (b) Find the value of $x^3 - 7x^2 - 2x + 88$ when $x = 5 + i\sqrt{3}$. 2

- 12.(a) Find the equation whose roots are squares of the differences of the roots of the equation $x^3 + x + 2 = 0$. 3
- (b) Transform the equation $x^4 + 4x^3 + 7x^2 + 6x - 4 = 0$ into one in which the terms involving x^3 is absent. 3

GROUP-C

Answer any two questions

12×2 = 24

- 13.(a) If $(x^2 + px + 1)$ be a factor of $(ax^3 + bx + c)$, then prove that $a^2 - c^2 = ab$. 4
- (b) If the equation $x^n - nqx + (n-1)r = 0$ has a pair of equal roots, show that $q^n = r^{n-1}$. 4
- (c) Show that if the equation $x^3 - ax^2 + bx - c = 0$ has a pair of roots of the form $\alpha(1 \pm i)$ where α is real, then $(a^2 - 2b)(b^2 - 2ac) = c^2$. 4
- 14.(a) If α, β, γ are the roots of the equation $x^3 - 3x^2 + x - 1 = 0$, then find the equation whose roots are

$$\alpha\beta + \frac{1}{\alpha} - \frac{1}{\beta}, \beta\gamma + \frac{1}{\beta} - \frac{1}{\gamma}, \gamma\alpha + \frac{1}{\gamma} - \frac{1}{\alpha}.$$
 6
- (b) If $\frac{p}{q}$ is a root of $a_0x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n = 0$, where $a_0, a_1, \dots, a_{n-1}, a_n$ are integers and p, q are integers prime to each other, then prove that q is a factor of a_0 and p is a factor of a_n . 6
- 15.(a) If $\alpha_1, \alpha_2, \dots, \alpha_n$ be the roots of $x^n + p_1x^{n-1} + \dots + p_{n-1}x + p_n = 0$, then find the value of $(\alpha_1^2 + 1)(\alpha_2^2 + 1) \dots (\alpha_n^2 + 1)$. 6
- (b) If α, β, γ are the roots of $x^3 + px + q = 0$, prove that $6S_5 = 5S_2S_3$, where $S_r = \sum \alpha^r$. 6
- 16.(a) Find the relation between a, b, c, d so that the product of two roots of the equation $x^4 + ax^3 + bx^2 + cx + d = 0$ is 1. 6
- (b) Show that the equation $(x-a)^3 + (x-b)^3 + (x-c)^3 + (x-d)^3 = 0$, where a, b, c, d are positive and not all equal, has only one real root. 6

—x—