

'समानों मन्त्रः समितिः समानी' UNIVERSITY OF NORTH BENGAL B.Sc. Honours 1st Semester Examination, 2023

CC1-MATHEMATICS

CALCULUS AND GEOMETRY

(REVISED SYLLABUS 2023)

Time Allotted: 2 Hours

Full Marks: 60

 $3 \times 4 = 12$

The figures in the margin indicate full marks.

GROUP-A

1. Answer any *four* questions from the following:

- (a) Determine the length of one arch of the cycloid $x = a(\theta \sin \theta)$, $y = a(1 \cos \theta)$.
- (b) Show that the curve $y = x^3$ has a point of inflexion at x = 0.

(c) Evaluate
$$\lim_{x\to 0} \frac{xe^x - \log(1+x)}{x^2}.$$

- (d) Find the points on the curve $y = x^4 6x^3 + 13x^2 10x + 5$, where the tangent is parallel to the line y = 2x.
- (e) Find the equation of the circle on the sphere $x^2 + y^2 + z^2 = 49$ whose centre is at the point (2, -1, 3).

(f) If
$$I_n = \int_{0}^{\pi/4} \tan^n x \, dx$$
, show that $I_{n+1} + I_{n-1} = \frac{1}{n}$

GROUP-B

- 2. Answer any *four* questions from the following: (a) If $y = \sin(m\cos^{-1}\sqrt{x})$, then prove that $\lim_{x \to 0} \frac{y_{n+1}}{y_n} = \frac{4n^2 - m^2}{4n+2}$. (b) Obvious the standard for the formula of $\frac{\pi/4}{4}$ is the standard formula of $\frac{\pi}{4}$ is the standard formula of \frac{\pi}{4} is the standard formula of $\frac{\pi}{4}$ is the standard formula of \frac{\pi}{4} is the standard formula of \frac{
 - (b) Obtain the reduction formula for $\int_{0}^{\pi/4} \sec^{n} x \, dx$ where $n \ (>1)$ being a positive 4+2 integer. Using this find the value of $\int_{0}^{\pi/4} \sec^{4} x \, dx$.

UG/CBCS/B.Sc./Hons./1st Sem./Mathematics/MATHCC1/Revised & Old/2023

(c) Transform the following equation to canonical form and determine the conic represented by it

$$x^2 + 4xy + 4y^2 - 20x + 10y - 50 = 0$$

- (d) Find the equation of the cylinder whose generators are parallel to the straight line 6 -3x = 6y = 2z and whose guiding curve is the ellipse $2x^2 + y^2 = 1$, z = 0.
- (e) *PSP'* is a focal chord of a conic $\frac{l}{r} = 1 + e \cos \theta$. Prove that the angle between the tangents at P and P' is $\tan^{-1}\left(\frac{2e\sin\alpha}{1-e^2}\right)$, where α is the angle between the chord and the major axis.
- (f) Show that the asymptotes of the curve $x^2y^2 = a^2(x^2 + y^2)$ form a square whose 6 side is of length 2a.

GROUP-C

3. $12 \times 2 = 24$ Answer any *two* questions from the following: (a) (i) Find the envelope of the family of ellipses $\frac{(x-\alpha)^2}{\alpha^2} + \frac{(y-\beta)^2}{b^2} = 1$, where

the parameters α and β are connected by $\frac{\alpha^2}{\alpha^2} + \frac{\beta^2}{b^2} = 1$.

- (ii) If $y = e^{m \sin^{-1} x}$, prove that $(1 x^2)y_{n+2} (2n+1)xy_{n+1} (m^2 + n^2)y_n = 0$. Obtain y_n for x = 0.
- (b) (i) Find the area in the first quadrant bounded by x = 0, y = 0 and 6+6 $\sqrt{x} + \sqrt{y} = \sqrt{a}$.
 - (ii) Find the volume of revolution generated by the region enclosed by $y = \sqrt{x}$ and the lines y = 1, x = 4 about x-axis.
- A tangent to the parabola $y^2 + 4bx = 0$ meets the parabola $y^2 = 4ax$ at P 6+6(c) (i) and Q. Prove that the locus of mid-point of PQ is $y^2(2a+b) = 4a^2x$.
 - (ii) Show that the distance between two points in two dimensional plane does not change under translation and rotation of co-ordinate axes.
- (d) (i) Find equations of the generating lines of the hyperboloid $\frac{x^2}{4} + \frac{y^2}{9} \frac{z^2}{16} = 1$ 6+6which passes through the point (2, 3, -4).
 - (ii) A sphere of constant radius 2a passes through the origin O and meets the axes in A, B, C. Show that the locus of the centroid of the tetrahedron *OABC* is the sphere $x^2 + y^2 + z^2 = a^2$.

_×__

6+6

6

6



UNIVERSITY OF NORTH BENGAL

B.Sc. Honours 1st Semester Examination, 2023

CC1-MATHEMATICS

CALCULUS, GEOMETRY AND DIFFERENTIAL EQUATION

(OLD SYLLABUS 2018)

Time Allotted: 2 Hours

Full Marks: 60

 $3 \times 4 = 12$

The figures in the margin indicate full marks.

GROUP-A

1. Answer any *four* questions from the following:

(a) Determine the length of one arch of the cycloid $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$.

(b) Show that the curve $y = x^3$ has a point of inflexion at x = 0.

(c) Evaluate
$$\lim_{x\to 0} \frac{xe^x - \log(1+x)}{x^2}.$$

(d) Solve
$$\log\left(\frac{dy}{dx}\right) = ax + by$$
.

(e) Find the equation of the circle on the sphere $x^2 + y^2 + z^2 = 49$ whose centre is at the point (2, -1, 3).

(f) If
$$I_n = \int_0^{\pi/4} \tan^n x \, dx$$
, show that $I_{n+1} + I_{n-1} = \frac{1}{n}$.

GROUP-B

2. Answer any *four* questions from the following: $6 \times 4 = 24$

(a) If
$$y = \sin(m\cos^{-1}\sqrt{x})$$
, prove that $\lim_{x \to 0} \frac{y_{n+1}}{y_n} = \frac{4n^2 - m^2}{4n+2}$. 6

(b) Obtain the reduction formula for
$$\int_{0}^{\pi/4} \sec^{n} x \, dx$$
 where $n(>1)$ being a positive $4+2$ integer. Using this find the value of $\int_{0}^{\pi/4} \sec^{4} x \, dx$.

3

UG/CBCS/B.Sc./Hons./1st Sem./Mathematics/MATHCC1/Revised & Old/2023

(c) Transform the following equation to its canonical form and determine the conic represented by it

$$x^2 + 4xy + 4y^2 - 20x + 10y - 50 = 0$$

(d) Find the equation of the cylinder whose generators are parallel to the straight line -3x = 6y = 2z and whose guiding curve is the ellipse $2x^2 + y^2 = 1$, z = 0.

6

 $12 \times 2 = 24$

6+6

(e) Solve
$$\frac{dy}{dx} = \sqrt{y-x}$$
.

(f) Show that the asymptotes of the curve $x^2y^2 = a^2(x^2 + y^2)$ form a square whose 6 side is of length 2a.

GROUP-C

3. Answer any *two* questions from the following:

(a) (i) Find the envelope of the family of ellipses $\frac{(x-\alpha)^2}{a^2} + \frac{(y-\beta)^2}{b^2} = 1$, where 6+6 the parameters α and β are connected by $\frac{\alpha^2}{a^2} + \frac{\beta^2}{b^2} = 1$.

(ii) If
$$y = e^{m \sin^{-1} x}$$
, prove that $(1 - x^2)y_{n+2} - (2n+1)xy_{n+1} - (m^2 + n^2)y_n = 0$.

- (b) (i) Find the area in the first quadrant bounded by x = 0, y = 0 and 6+6 $\sqrt{x} + \sqrt{y} = \sqrt{a}$.
 - (ii) Find the volume of revolution generated by the region enclosed by $y = \sqrt{x}$ and the lines y = 1, x = 4 about x-axis.

(c) (i) Solve
$$\tan y \frac{dy}{dx} = \sin(x+y) + \sin(x-y)$$
.

- (ii) Solve $\frac{dy}{dx} + 2xy = xy^3$.
- (d) (i) Find equations of the generating lines of the hyperboloid $\frac{x^2}{4} + \frac{y^2}{9} \frac{z^2}{16} = 1$. 6+(3+3) which passes through the point (2, 3, -4).
 - (ii) Show that the distance between two points in two dimensional plane does not change under translation and rotation of co-ordinate axes.

×