



‘समानो मन्त्रः समितिः समानी’

UNIVERSITY OF NORTH BENGAL

B.Sc. Honours 1st Semester Examination, 2021

GE1-P1-MATHEMATICS

Time Allotted: 2 Hours

Full Marks: 60

*The figures in the margin indicate full marks.
All symbols are of usual significance.*

**The question paper contains GE1, GE2, GE3, GE4 and GE5.
Candidates are required to answer any *one* from the *five* courses and
they should mention it clearly on the Answer Book.**

GE1

CALCULUS, GEOMETRY AND DIFFERENTIAL EQUATION

GROUP-A

1. Answer any **four** questions from the following: 3×4 = 12
 - (a) Find the points of inflexion on the curve $(\theta^2 - 1)r = a\theta^2$. 3
 - (b) Find the envelopes of the lines $\frac{x}{a} + \frac{y}{b} = 1$, where a and b are parameters related by $a + b = c$. 3
 - (c) Find the equation of the sphere through the circle $x^2 + y^2 + z^2 = 9$, $x + y - 2z = 4$ and the origin. 3
 - (d) Evaluate $\int_0^1 xe^{-\sqrt{x}} dx$ using reduction formula. 3
 - (e) Obtain the singular solution of the equation $(xp - y)^2 = p^2 - 1$, where $p = dy/dx$. 3
 - (f) Determine the nature of the quadric $5x^2 - 6xy + 5y^2 + 22x - 26y + 29 = 0$. 3

GROUP-B

2. Answer any **four** questions from the following: 6×4 = 24
 - (a) If $y = \frac{\sin^{-1} x}{\sqrt{1-x}}$, $|x| < 1$, then prove that $(1-x^2)y_{n+2} - (2n+3)xy_{n+1} - (n+1)^2 y_n = 0$. 6
 - (b) Find the asymptotes of the curve $x^3 + 2x^2y - 4xy^2 + 8y^3 - 4x + 8y - 10 = 0$. 6
 - (c) Find the area of the region lying between the cissoid $y^2 = \frac{x^3}{2a-x}$ and its asymptote. 6
 - (d) Solve: $y(2xy + 1) dx + x(1 + 2xy + x^2y^2) dy = 0$ 6
 - (e) Find a, b such that, $\lim_{x \rightarrow 0} \frac{x(1 + a \cos x) - b \sin x}{x^3} = 1$. 6
 - (f) Find the volume of the solid generated by revolving the cardioid $r = a(1 + \cos \theta)$ about initial line. 6

GROUP-C

Answer any *two* questions from the following

12×2 =24

3. (a) Find the range of values of x for which $y = x^4 - 6x^3 + 12x^2 + 5x + 7$ is concave upward or downward. 6
 (b) Find the length of the arc of the cardioid $r = a(1 - \cos \theta)$ lying inside the circle $r = a \cos \theta$. 6
4. (a) Solve by using Bernoulli form $\frac{dy}{dx} + \frac{y}{x} \log y = \frac{y}{x^2} (\log y)^2$. 7
 (b) Solve: $(xy^2 - e^{1/x^3}) dx - x^2 y dy = 0$ 5
5. (a) Reduce the equation $7x^2 - 2xy + 7y^2 - 16x + 16y - 8 = 0$ to its canonical form and hence determine the nature of the conic. 8
 (b) Find the equation of the sphere for which the circle $x^2 + y^2 + z^2 + 7y - 2z + 2 = 0$, $2x + 3y + 4z = 8$ is a great circle. 4
6. (a) Find the value of $y_n(0)$, where $y = \log(x + \sqrt{1 + x^2})$. 6
 (b) If $I_{m,n} = \int_0^{\pi/2} \cos^m x \sin nx dx$, then show that $I_{m,n} = \frac{1}{m+n} + \frac{m}{m+n} I_{m-1, n-1}$. 6

GE2

ALGEBRA

GROUP-A

1. Answer any *four* questions from the following: 3×4 = 12
 (a) Apply Descartes's rule of signs to find the nature of the roots of the equation $x^4 + mx^2 + nx - p = 0$, where m, n, p are positive. 3
 (b) Prove that $\sqrt{i} + \sqrt{-i} = \sqrt{2}$. 3
 (c) Prove that the eigenvalues of a real skew symmetric matrix are purely imaginary or zero. 3
 (d) Find the sum of 99th power of the roots of the equation $x^7 - 1 = 0$. 3
 (e) Use Cayley-Hamilton theorem to find A^{-1} for the matrix 3

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 2 & 1 \\ 2 & 3 & 2 \end{bmatrix}$$

 (f) Find the quadratic equation whose roots are twice the roots of $2x^2 - 5x + 2 = 0$. 3

GROUP-B

2. Answer any *four* questions from the following: 6×4 = 24
 (a) If $2 \cos \theta = x + \frac{1}{x}$ and θ is real, prove that $2 \cos n\theta = x^n + \frac{1}{x^n}$, n being an integer. 6

- (b) Solve the equation $16x^4 - 64x^3 + 56x^2 + 16x - 15 = 0$ whose roots are in arithmetic progression. 6
- (c) Find integers u and v satisfying $52u - 91v = 78$. 6
- (d) Find all eigenvalues and eigenvectors of the matrix $A = \begin{bmatrix} 2 & 0 & 1 \\ 4 & 3 & 1 \\ 2 & 1 & 1 \end{bmatrix}$. 6
- (e) For what values of λ the following system of equations are consistent? 6
- $$\begin{aligned} x - y + z &= 1 \\ x + 2y + 4z &= \lambda \\ x + 4y + 6z &= \lambda^2 \end{aligned}$$
- (f) Use Cayley-Hamilton theorem to find A^{100} , where $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$. 6

GROUP-C

Answer any two questions from the following

12×2= 24

3. (a) If $\log \sin(\theta + i\phi) = \alpha + i\beta$, then prove that $2e^{2\alpha} = \cosh 2\phi - \cos 2\theta$. 6
- (b) Find the relation among the coefficients of the equation $a_0x^4 + 4a_1x^3 + 6a_2x^2 + 4a_3x + a_4 = 0$, so that the second term and the fourth term may be removed by the transformation $x = y + h$. 6
4. (a) If α, β, γ are the roots of the equation $x^3 + qx + r = 0$, find the equation whose roots are $\beta + \gamma - 2\alpha, \gamma + \alpha - 2\beta, \alpha + \beta - 2\gamma$. 6
- (b) Determine all values of $(1 + i\sqrt{3})^{3/4}$ and show that their product is 8. 4+2
5. (a) Solve the equation $3x^3 + 5x^2 + 5x + 3 = 0$, which has three distinct roots of equal moduli. 6
- (b) If roots of $ax^3 + bx^2 + cx + d = 0$ are in arithmetic progression. Show that $2b^3 - 9abc + 27a^2d = 0$. 6
6. (a) Determine the conditions for which the system of equation has 2+2+2
- (i) only one solution
 - (ii) no solution
 - (iii) infinitely many solution.
- $$\begin{aligned} x + 2y + z &= 1 \\ 2x + y + 3z &= b \\ x + ay + 3z &= b + 1 \end{aligned}$$
- (b) The matrix of a linear mapping $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ with ordered basis $\{(0, 1, 1), (1, 0, 1), (1, 1, 0)\}$ of \mathbb{R}^3 is given by 6
- $$\begin{pmatrix} 0 & 3 & 0 \\ 2 & 3 & -2 \\ 2 & -1 & 2 \end{pmatrix}$$
- Find the matrix of T relative to the ordered basis $\{(2, 1, 1), (1, 2, 1), (1, 1, 2)\}$ of \mathbb{R}^3 .

GE3

DIFFERENTIAL EQUATION AND VECTOR CALCULUS

GROUP-A

1. Answer any **four** questions from the following: 3×4 = 12
- (a) Show that the function $f(x, y) = xy^2$ does not satisfy the Lipschitz condition on the strip $|x| \leq 1, |y| < \infty$.
- (b) Find the Wronskian of $\{1, 1+x, 1+x+x^2+x^3\}$.
- (c) Define Lipschitz constant. Find Lipschitz constant for the function $f(x, y) = x^2y^2$ defined on $|x| \leq 1, |y| \leq 1$.
- (d) Solve: $\frac{d^5y}{dx^5} - 2\frac{d^4y}{dx^4} + \frac{d^3y}{dx^3} = 0$
- (e) Examine whether the vector valued function $\vec{r} = t^3\hat{i} + e^t\hat{j} + \frac{1}{t+3}\hat{k}$ is continuous at $t = -3$ or not.
- (f) Evaluate: $\lim_{t \rightarrow 1} \left[\frac{t^3-1}{t-1}\hat{i} + \frac{t^2-3t+2}{t^2+t-2}\hat{j} + (t^2+1)e^{t-1}\hat{k} \right]$

GROUP-B

2. Answer any **four** questions from the following: 6×4 = 24
- (a) (i) If y_1 and y_2 are two independent solutions of the linear equation $\frac{d^2y}{dx^2} + p\frac{dy}{dx} + qy = 0$, then show that the Wronskian $W(y_1, y_2) = Ae^{-\int p dx}$, where A is a constant. 3+3
- (ii) Show that the functions $\{e^{2x}, e^{2x} \cos 4x, e^{2x} \sin 4x\}$ are linearly independent.
- (b) Show that linearly independent solutions of $y'' - 2y' + 2y = 0$ are $e^x \sin x$ and $e^x \cos x$. What is the general solution? Find the solution $y(x)$ with the conditions $y(0) = 2, y'(0) = -3$. 6
- (c) Solve: $x^3 \frac{d^3y}{dx^3} + 2x^2 \frac{d^2y}{dx^2} + 2y = 10(x+x^{-1})$ 6
- (d) Solve: $(D^3 - 1)y = x \sin x, D \equiv \frac{d}{dx}$ 6
- (e) (i) Find the co-ordinates of the point where the line $\vec{r} = t\hat{i} + (1+2t)\hat{j} - 3t\hat{k}$ intersects the plane $3x - y - z = 2$. 3+3
- (ii) Show that the graph of $\vec{r}(t) = t\hat{i} + \frac{1+t}{t}\hat{j} + \frac{1-t^2}{t}\hat{k}, t > 0$ lies on the plane $x - y + z + 1 = 0$.
- (f) (i) Find the domain of the vector function $h(t)F(t)$, where $h(t) = \sin t$ and $F(t) = \frac{1}{\cos t}\hat{i} + \frac{1}{\sin t}\hat{j} + \frac{1}{\tan t}\hat{k}$. 3+3
- (ii) Find $(F \times G)(t)$ if $F(t) = t^2\hat{i} + t\hat{j} - (\sin t)\hat{k}$ and $G(t) = t^2\hat{i} + \frac{1}{t}\hat{j} + 5\hat{k}$.

GROUP-C

Answer any *two* questions from the following

12×2 =24

3. (a) (i) Solve by the method of variation of parameters $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + y = xe^x \log x$. 5+3+4
- (ii) Evaluate: $\frac{1}{D^2 - 3D + 2} xe^{3x}$
- (iii) Solve: $\frac{d^2y}{dx^2} + 16y = 1$, $y(0) = 1$, $y'(0) = 2$
- (b) (i) Solve by using the method of undetermined coefficient 6+6
 $(D^2 + D - 6)y = 10e^{2x} - 18e^{3x} - 6x - 11$
- (ii) Solve: $(D^4 + 2D^3 - 3D^2)y = x^2 + 3e^{2x} + 4\sin x$
- (c) (i) Solve the equations 6+6

$$\begin{cases} \frac{dx}{dt} = -wy \\ \frac{dy}{dt} = wx \end{cases}$$

and show that the point (x, y) lies on a circle.

- (ii) Solve the system of equations

$$\begin{cases} \frac{dx}{dt} = -x + 6y \\ \frac{dy}{dt} = x - 2y \end{cases}$$

- (d) (i) Find the slope of the line in R^2 for the vector equation 4+1+4+3
 $\vec{r}(t) = (1 - 2t)\hat{i} - (2 - 5t)\hat{j}$
- (ii) Define continuity of a vector valued function.
- (iii) Show that the vector function $\vec{r}(t) = \begin{cases} \frac{\sin t}{t}\hat{i} + t\hat{j} + t^2\hat{k} & , t \neq 0 \\ \hat{i} & , t = 0 \end{cases}$
 is continuous at $t = 0$.
- (iv) Find a vector function F whose graph is the curve of intersection of the hemisphere $z = \sqrt{4 - x^2 - y^2}$ and the curve $y = x^2$.

GE4

GROUP THEORY

GROUP-A

1. Answer any *four* questions from the following: 3×4= 12
- (a) Let (S, \circ) be a semigroup. If for all $x, y \in S$, $x^2 \circ y = y \circ x^2$, prove that (S, \circ) is an abelian group. 3
- (b) Suppose H, K are subgroups of index 2 in a group G . Prove that $H \cap K$ is a normal subgroup of G . 3
- (c) Let $G = \langle a \rangle$ be a cyclic group of order n . Prove that every subgroup of G is of the form $\langle a^m \rangle$, where m is a divisor of n . 3
- (d) Find all elements of order 10 in the group $(\mathbb{Z}_{30}, +)$. 3

- (e) Show that there does not exist an onto homomorphism from the group $(\mathbb{Z}_6, +)$ to $(\mathbb{Z}_4, +)$. 3
- (f) Prove or disprove: $(\mathbb{Q}, +)$ is isomorphic to (\mathbb{Q}^+, \cdot) 3

GROUP-B

2. Answer any **four** questions from the following: 6×4 = 24
- (a) Let S be the set of all permutations on the set $\{1, 2, 3\}$. Show that S forms a non-abelian group with respect to multiplication. 6
- (b) Suppose that the order of an element a in a group (G, \circ) is n . Show that $O(a^m) = \frac{n}{d}$, where $d = \gcd(m, n)$. Find the order of $\overline{n-1}$ in $(\mathbb{Z}_n, +)$. 4+2
- (c) (i) Let H be a subgroup of a group G and $a, b \in G$. Prove that $b \in Ha$ iff $ba^{-1} \in H$. 2+4
- (ii) Let H be a subgroup of a group G . Show that the set of all distinct left cosets of H in G and the set of all distinct right cosets of H in G have the same cardinality.
- (d) (i) If H is a subgroup of G and N is a normal subgroup of G , then show that $H \cap N$ is a normal subgroup of H . 4+2
- (ii) Prove that N is a normal subgroup of G iff $gNg^{-1} = N$ for every $g \in G$.
- (e) (i) Let (G, \circ) be a group and H, K be subgroups of (G, \circ) . Show that HK forms a subgroup of (G, \circ) iff $HK = KH$. 4+2
- (ii) Check whether the union of two subgroups of a group (G, \circ) is a subgroup of (G, \circ) or not?
- (f) (i) Let (G, \circ) be a group and a mapping $\varphi: G \rightarrow G$ is defined by $\varphi(x) = x^2, x \in G$. Prove that φ is a homomorphism iff G is commutative. 4+2
- (ii) Prove that $(\mathbb{Z}_4, +)$ and “Klein’s 4-group” are not isomorphic.

GROUP-C

Answer any **two** questions from the following

12×2 =24

3. (a) (i) Let $G = S_3, G' = (\{-1, 1\}, \cdot)$ and $\varphi: G \rightarrow G'$ is defined by 4+2+2
- $$\varphi(\alpha) = \begin{cases} 1 & , \alpha \text{ be an even permutation in } S_3 \\ -1 & , \alpha \text{ be an odd permutation in } S_3 \end{cases}$$
- Then, (I) Show that φ is homomorphism.
 (II) Find $\ker \varphi$.
 (III) Deduce that A_3 is a normal subgroup of S_3 .
- (ii) Prove that a finite cyclic group of order n is isomorphic to $(\mathbb{Z}_n, +)$. 4
- (b) (i) Let H be a normal subgroup of G . Prove that the quotient group G/H is abelian iff $xyx^{-1}y^{-1} \in H$ for all $x, y \in G$. 4+4+4
- (ii) Suppose that a subgroup H of a group G has the property that $x^2 \in H$ for every $x \in G$. Prove that H is normal in G and G/H is abelian.
- (iii) Let $G = \left\{ \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} : a, b, d \in \mathbb{R} \text{ and } ad \neq 0 \right\}$ and $H = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} : b \in \mathbb{R} \right\}$. Show that H is a normal subgroup of G .

- (c) Let M be the set of all real matrices $\left\{ \begin{pmatrix} a & a \\ b & b \end{pmatrix} : a + b \neq 0 \right\}$. Prove that 4+4+4
- (i) (M, \circ) is a semi-group under matrix multiplication.
- (ii) there is no left identity in the semi-group.
- (iii) $\begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ is a right identity.
- (d) (i) Let $(G, \circ), (G', *)$ be two groups and $\varphi: (G, \circ) \rightarrow (G', *)$ be an onto homomorphism. Then prove that $G/\ker \varphi \cong G'$. 6+6
- (ii) Let G be a cyclic group of order 10 and G' be a cyclic group of order 5. Show that there exists a homomorphism φ of G onto G' with $o(\ker \varphi) = 2$.

GE5

NUMERICAL METHODS

GROUP-A

1. Answer any **four** questions from the following: 3×4 = 12
- (a) If $f(x) = 4 \cos x - 6x$, find the relative percentage error in $f(x)$ for $x = 0$, if the error in $x = 0.005$. 3
- (b) Deduce the iterative procedure $x_{n+1} = \frac{1}{2} \left(x_n + \frac{a}{x_n} \right)$ for evaluating \sqrt{a} using Newton-Raphson method. 3
- (c) Prove that $\left(\frac{\Delta^2}{E} \right) x^3 = 6xh^2$ where the notations used have their usual meanings. 3
- (d) Show that $\nabla y_{n+1} = h \left[1 + \frac{1}{2} \nabla + \frac{5}{12} \nabla^2 + \dots \right] Dy_n$, where D is the differential operator. 3
- (e) Write down the convergence of bisection method. 3
- (f) What is the geometrical significance of Simpson's one-third rule? 3

GROUP-B

2. Answer any **four** questions from the following: 6×4 = 24
- (a) If a number is connected to n significant figures and the first significant figure of the number is k , then prove that the relative error $\epsilon_r < \frac{1}{k \cdot 10^{n-1}}$. 6
- (b) Find the positive root of the equation $x^3 + x - 1 = 0$ by fixed point iteration method correct upto three decimal places. 6
- (c) Find a real root of the equation $x^x + 2x - 2 = 0$ correct upto five decimal places using bisection method. 6
- (d) Define backward difference operator ∇ and shifting operator E . Show that 6

$$\sum_{k=0}^{n-1} \Delta^2 f_k = \Delta f_n - \Delta f_0$$

- (e) Use Runge-Kutta method of order two to find $y(0.1)$ and $y(0.2)$ correct upto four decimal places given $\frac{dy}{dx} = y - x$, $y(0) = 2$. 6
- (f) Explain Gauss-Seidel method for solving a system of linear equations. Obtain the sufficient condition for convergence of Gauss-Seidel method. 6

GROUP-C

Answer any two questions from the following

12×2 =24

3. (a) Evaluate $\int_0^{\pi/2} \sqrt{1-0.162 \sin^2 \theta} d\theta$, by Simpson's $\frac{1}{3}$ rd rule, correct upto 4 decimal places taking 12 points. 6

- (b) Given $\frac{dy}{dx} = \frac{-y}{1+x}$, $y(0.3) = 2$. Compute $y(1)$ by Euler's method, correct upto four decimal places, taking step length $h = 0.1$. 6

4. (a) Solve the system of equations by Gauss-elimination method 6

$$\begin{aligned} 3x + 9y - 2z &= 11 \\ 4x + 2y + 13z &= 24 \\ 4x - 2y + z &= -8 \end{aligned}$$

correct upto 2 decimal places.

- (b) Using Newton-Raphson method find a positive root of the equation $e^x - 3x = 0$ correct upto four decimal places. 6

5. (a) Find $f(x)$ as a polynomial in x by using the following table: 6

x	0	2	4	6	8
$f(x)$	2.51881	2.53148	2.54407	2.55630	2.56820

- (b) Obtain the missing terms in the following table: 6

x	1	2	3	4	5	6	7	8
$f(x)$	1	8	*	64	*	216	343	512

6. (a) Explain the method of fixed point iteration with the condition of convergence for numerical solution of an equation of the form $x = \phi(x)$. 6

- (b) What is interpolation? Establish Lagrange's polynomial interpolation formula. 6

—x—